

COMBINATORIAL AND MODEL-THEORETICAL PRINCIPLES RELATED TO REGULARITY OF ULTRAFILTERS AND COMPACTNESS OF TOPOLOGICAL SPACES. VI.

PAOLO LIPPARINI

ABSTRACT. We discuss the existence of complete accumulation points of sequences in products of topological spaces. Then we collect and generalize many of the results proved in Parts I, II and IV.

The present Part VI is complementary to Part V to the effect that here we deal, say, with uniformity, complete accumulation points and κ -(λ)-compactness, rather than with regularity, $[\lambda, \mu]$ -compactness and κ -(λ, μ)-compactness. Of course, if we restrict ourselves to regular cardinals, Parts V (for $\lambda = \mu$) and Part VI essentially coincide.

See Parts I - IV [L7], or [BF, C2, CK, CN, KM, KV, L1, L2, L3, L4, L5, L6, S, V1] for unexplained notation.

Let us recall the definition of the $\square^{<\nu}$ product. If ν is a cardinal, and $(X_\beta)_{\beta \in \kappa}$ is a family of topological spaces, then the $\square^{<\nu}$ topology on the cartesian product $\prod_{\beta \in \kappa} X_\beta$ is the topology a base of which is given by all products $\prod_{\beta \in \kappa} Y_\beta$, where each Y_β is an open subset of X_β , and $|\{\beta \in \kappa \mid Y_\beta \neq X_\beta\}| < \nu$. The product of $(X_\beta)_{\beta \in \kappa}$ with the $\square^{<\nu}$ topology shall be denoted by $\square_{\beta \in \kappa}^{<\nu} X_\beta$.

Notice that in the case $\nu = \omega$ we get the more usual Tychonoff product. As usual, we shall denote the Tychonoff product by $\prod_{\beta \in \kappa} X_\beta$.

Recall that, for every infinite cardinal λ , a topological space X is said to satisfy CAP_λ if and only if every subset $Y \subseteq X$ with $|Y| = \lambda$ has a complete accumulation point in X .

2000 *Mathematics Subject Classification.* Primary 03C20, 54D20, 03E05, 03C95, 03E75; Secondary 54B10, 54A35, 54F05, 03C55, 03C98.

Key words and phrases. Complete accumulation points, CAP_λ , compactness of products of topological spaces; ν -complete, uniform, regular, μ -decomposable ultrafilters; elementary extensions of cardinals with additional structure; μ -non standard elements; infinite matrices combinatorics; compactness of abstract logics.

The author has received support from MPI and GNSAGA. We wish to express our gratitude to X. Caicedo for stimulating discussions and correspondence.

Actually, in what follows we shall use the variation of CAP_λ which deals with sequences, rather than subsets.

If λ is an infinite cardinal, X is a topological space and $(x_\alpha)_{\alpha < \lambda}$ is a sequence of elements of X , we say that $x \in X$ is a λ -*accumulation point* for $(x_\alpha)_{\alpha < \lambda}$ if and only if $|\{\alpha < \lambda \mid x_\alpha \in U\}| = \lambda$, for every neighborhood U of x .

We say that a topological space X satisfies CAP_λ^* if and only if every λ -indexed sequence $(x_\alpha)_{\alpha < \lambda}$ of elements of X has a λ -accumulation point.

Proposition 1. (a) *If λ is a regular cardinal, then CAP_λ^* is equivalent to CAP_λ , and equivalent to $[\lambda, \lambda]$ -compactness.*

(b) *If λ is a singular cardinal, then CAP_λ^* is equivalent to the conjunction of CAP_λ and $\text{CAP}_{\text{cf } \lambda}$.*

Notice that the space $\text{cf } \lambda$, endowed with the order topology, does not satisfy $\text{CAP}_{\text{cf } \lambda}$, but it satisfies CAP_κ , for every infinite cardinal $\kappa \neq \text{cf } \lambda$.

On the other hand, if μ is regular, and $\mu \leq \lambda$, consider the space $X = S_\mu(\lambda)$, endowed with the topology which has, as a base, all the sets of the form $\{y \in S_\mu(\lambda) \mid y \subseteq x\}$, x varying in $S_\mu(\lambda)$. X satisfies CAP_κ , for every infinite cardinal $\kappa < \mu$. Indeed, if $\kappa < \mu$, then every κ -indexed sequence $(x_\beta)_{\beta \in \kappa}$ in X converges to $\bigcup_{\beta \in \kappa} x_\beta$. On the contrary, if $\lambda \geq \kappa \geq \mu$, then X fails to satisfy CAP_κ .

Thus, if in the above example we take λ singular, and $\mu = (\text{cf } \lambda)^+$, we get a space satisfying $\text{CAP}_{\text{cf } \lambda}$ but not CAP_λ .

If λ is singular, and $2^{\text{cf } \lambda} < \lambda$, then it is not difficult to construct a Tychonoff topological space X which does not satisfy CAP_λ , but such that every power of X satisfies $\text{CAP}_{\text{cf } \lambda}$.

Notice that if λ is singular, $\text{CAP}_{\text{cf } \lambda}$ holds, and there are arbitrarily large $\kappa < \lambda$ such that CAP_κ holds, then CAP_λ holds, hence, by Proposition 1(b), also CAP_λ^* holds. The proof of the above fact is similar to the proof of the proposition in [S, p. 94].

Proposition 2. *Suppose that X is a topological space, $x \in X$, λ is an infinite cardinal and $(x_\gamma)_{\gamma \in \lambda}$ is a sequence of elements of X .*

Then x is a λ -accumulation point for $(x_\gamma)_{\gamma \in \lambda}$ if and only if there exists some ultrafilter D uniform over λ such that $(x_\gamma)_{\gamma \in \lambda}$ D -converges to x .

Remark 3. Variations on the above proposition are well known. See e.g. [S, p. 80-81].

An analogue of the above proposition for (λ', λ) -regular ultrafilters and $[\lambda, \lambda']$ -compactness is proved in [C1, C2].

For $\lambda = \lambda'$ a regular cardinal, the statement in Proposition 2 and the corresponding result from [C1, C2] essentially amount to the same result, since an ultrafilter is (λ, λ) -regular if and only if it has some quotient uniform over λ , and since CAP_λ^* is equivalent to $[\lambda, \lambda]$ -compactness.

Notations 4. Since we will often be working in products, dealing with sequences indexed by external sets, in order to avoid confusion we shall introduce a special notation to denote the elements of the product.

If $x \in \prod_{\beta \in \kappa} X_\beta$, say $x = (x_\beta)_{\beta \in \kappa}$, we shall sometimes denote x by $\prod_{\beta \in \kappa} x_\beta$.

The next lemma generalizes the fundamental property of D -convergence with respect to products. Indeed, since every ultrafilter is ω -complete, the particular case $\nu = \omega$ of the next lemma asserts that D -convergence in a Tychonoff product is equivalent to factor by factor D -convergence. The lemma is more general in the sense that it asserts the above equivalence for ν -complete ultrafilters and $\square^{<\nu}$ products.

Lemma 5. *Suppose that D is a ν -complete ultrafilter over some set I , $(X_\beta)_{\beta \in \kappa}$ is a family of topological spaces, and $(x_i)_{i \in I}$ is a sequence of elements of $X = \square_{\beta \in \kappa}^{<\nu} X_\beta$, say, for each $i \in I$, $x_i = \prod_{\beta \in \kappa} x_{i,\beta}$.*

Then $(x_i)_{i \in I}$ D -converges in X to some $x = \prod_{\beta \in \kappa} x_\beta \in X$ if and only if, for each $\beta \in \kappa$, $(x_{i,\beta})_{i \in I}$ D -converges to x_β in X_β .

Proposition 6. *Suppose that κ is a cardinal, and $\lambda, \nu, (\mu_\beta)_{\beta \in \kappa}$ are infinite cardinals. For each $\beta \in \kappa$, endow μ_β with the order topology. Suppose that $(f_\beta)_{\beta \in \kappa}$ is a given set of functions such that $f_\beta : \lambda \rightarrow \mu_\beta$, for each $\beta \in \kappa$.*

Consider the following conditions.

- (a) *There exists a ν -complete ultrafilter D uniform over λ such that for no $\beta \in \kappa$ $f_\beta(D)$ is uniform over μ_β .*
- (b) *In the space $\square_{\beta \in \kappa}^{<\nu} \mu_\beta$ the sequence $(x_\gamma)_{\gamma < \lambda}$ defined by $x_\gamma = \prod_{\beta \in \kappa} f_\beta(\gamma)$ has a λ -accumulation point in $\square_{\beta \in \kappa}^{<\nu} \mu_\beta$.*

If for each $\beta \in \kappa$ μ_β is a regular cardinal, then Condition (a) implies Condition (b).

If $\nu = \omega$ then Condition (b) implies Condition (a).

Proof. (a) $\Rightarrow$ (b). Fix $\beta \in \kappa$. Since $f_\beta(D)$ is not uniform over μ_β , and μ_β is a regular cardinal, then $f_\beta(D)$ is not (μ_β, μ_β) -regular. By [L5, Proposition 1], μ_β is $f_\beta(D)$ -compact, that is, every $f_\beta(D)$ -sequence in μ_β $f_\beta(D)$ -converges. In particular, the identity function on $f_\beta(\lambda)$ $f_\beta(D)$ -converges to some point of μ_β . By definition of $f_\beta(D)$ and of ultrafilter convergence, this implies that the sequence $(f_\beta(\gamma))_{\gamma < \lambda}$ D -converges in μ_β to some point, say, to x_β .

Since the above holds for every $\beta \in \kappa$, and $(x_\gamma)_\beta = f_\beta(\gamma)$, then, by Lemma 5, the sequence $(x_\gamma)_{\gamma < \lambda}$ D -converges in $\square_{\beta \in \kappa}^{\leq \nu} \mu_\beta$ to $x = \prod_{\beta \in \kappa} x_\beta$. By Proposition 2, x is a λ -accumulation point for $(x_\gamma)_{\gamma < \lambda}$.

Now suppose that $\nu = \omega$. We are going to show that (b) implies (a). Suppose that (b) holds. By Proposition 2 there exists some ultrafilter D uniform over λ such that $(x_\gamma)_{\gamma < \lambda}$ D -converges to some point of $\prod_{\beta \in \kappa} \mu_\beta$.

By Lemma 5, for each $\beta \in \kappa$, the sequence $((x_\gamma)_\beta)_{\gamma < \lambda}$ D -converges to some point of μ_β . Since $(x_\gamma)_\beta = f_\beta(\gamma)$, this implies that, for each $\beta \in \kappa$, the identity function on μ_β $f_\beta(D)$ -converges to some point of μ_β , and this easily implies that $f_\beta(D)$ is not uniform over μ_β . $\square$

Definition 7. Suppose that λ is an infinite cardinal, and $\mathfrak{M}$ is a model with a unary predicate U and a distinguished binary relation $<$ such that $\langle U^{\mathfrak{M}}, <^{\mathfrak{M}} \rangle = \langle \lambda, < \rangle$. Suppose further that $\mathfrak{M}$ has a name for every element of $U = \lambda$ (for simplicity, and by abuse of notation, we shall suppose that the name for α is α itself).

If $\mathfrak{N} \equiv \mathfrak{M}$, and $b \in N$, we say that b is λ -non standard if and only if, in $\mathfrak{N}$, $U(b)$ holds, and $\alpha < b$ holds for every $\alpha < \lambda$. Other sentences which have been used to indicate similar notions are that “ b bounds λ ”, or that “ b realizes λ ”. Of course, in the particular case $\lambda = \omega$, we get the usual notion of a non-standard element. In other words, $\mathfrak{N}$ has a λ -non standard element if and only if $\langle U^{\mathfrak{N}}, <^{\mathfrak{N}} \rangle$ is not an *end extension* of $\langle \lambda, < \rangle$.

We shall use the above terminology even when U is not a predicate in the vocabulary of $\mathfrak{M}$, but just a unary relation definable by some formula. In particular, if $\mu < \lambda$, and, as above, in $\mathfrak{M}$, $\langle U, < \rangle = \langle \lambda, < \rangle$, $\mathfrak{M}$ has a name for every element of λ , and $\mathfrak{N} \equiv \mathfrak{M}$, we shall say that b is μ -non standard in $\mathfrak{N}$ if and only if, in $\mathfrak{N}$, $b < \mu$ holds, and $\alpha < b$ holds for every $\alpha < \mu$.

Of course, it might be the case that the model $\mathfrak{M}$ has many predicates U_i and relations R_i such that $\langle U_i, R_i \rangle \cong \langle \lambda, < \rangle$. If this is the case, and $\mathfrak{N} \equiv \mathfrak{M}$, it might happen that $\mathfrak{N}$ has a λ -non standard element according to, say, U_0, R_0 , but no λ -non standard element according to U_1, R_1 . We shall try to adhere to the convention that the base set of $\mathfrak{M}$ does always contain the “real” λ , so that our definition of λ -non standard is not ambiguous. Anyway, the above possible ambiguity is not a serious problem, as far as the present paper is concerned, since we always allow $\mathfrak{M}$ to be expanded with additional functions and relations, so that we can always have, inside $\mathfrak{M}$, isomorphisms between any $\langle U_i, R_i \rangle$ and any $\langle U_j, R_j \rangle$. Since being an isomorphism is expressible by first order sentences, and $\mathfrak{N}$ is supposed to be elementarily equivalent

to $\mathfrak{M}$, then a λ -non standard element exists in $\mathfrak{N}$ according to U_i, R_i if and only if a λ -non standard element exists according to U_j, R_j .

As far as Condition (7) below is concerned, fix some set $V \subseteq S_\lambda(\lambda)$ cofinal in $S_\lambda(\lambda)$ of cardinality $\leq \kappa$ and, for $v \in V$, let R_v be the unary predicate on λ defined by $R_v(\gamma)$ if and only if $\gamma \in v$ (compare Part IV, Definition 2 and Remark 3).

As far as Condition (8) below is concerned, recall the definition of a κ -(λ)-compact logic from Part IV, Definition 10.

Theorem 8. *Suppose that κ is a cardinal, λ is an infinite cardinal, and $(\mu_\beta)_{\beta \in \kappa}$ is a set of infinite regular cardinals.*

Then the following conditions are equivalent.

- (1) *There are κ functions $(f_\beta)_{\beta \in \kappa}$ such that*
 - (a) *For each $\beta \in \kappa$, $f_\beta : \lambda \rightarrow \mu_\beta$; and*
 - (b) *for every ultrafilter D uniform over λ there is $\beta \in \kappa$ such that $f_\beta(D)$ is uniform over μ_β .*
- (2) *There are κ functions $(f_\beta)_{\beta \in \kappa}$ such that*
 - (a) *For each $\beta \in \kappa$, $f_\beta : \lambda \rightarrow \mu_\beta$; and*
 - (b) *for every function $g \in \prod_{\beta \in \kappa} \mu_\beta$ there exists some finite set $F \subseteq \kappa$ such that $\left| \bigcap_{\beta \in F} f_\beta^{-1}([0, g(\beta))) \right| < \lambda$.*
- (3) *There is a family $(B_{\alpha, \beta})_{\beta \in \kappa, \alpha < \mu_\beta}$ of subsets of λ such that:*
 - (a) *For every $\beta \in \kappa$, $\bigcup_{\alpha < \mu_\beta} B_{\alpha, \beta} = \lambda$;*
 - (b) *For every $\beta \in \kappa$ and $\alpha \leq \alpha' < \mu_\beta$, $B_{\alpha, \beta} \subseteq B_{\alpha', \beta}$;*
 - (c) *For every function $g \in \prod_{\beta \in \kappa} \mu_\beta$ there exists a finite set $F \subseteq \kappa$ such that $\left| \bigcap_{\beta \in F} B_{g(\beta), \beta} \right| < \lambda$.*
- (4) *For every family $(X_\beta)_{\beta \in \kappa}$ of topological spaces, if $\prod_{\beta \in \kappa} X_\beta$ satisfies CAP_λ^* , then there is $\beta \in \kappa$ such that X_β satisfies $\text{CAP}_{\mu_\beta}^*$.*
- (5) *The topological space $\prod_{\beta \in \kappa} \mu_\beta$ does not satisfies CAP_λ^* , where each μ_β is endowed with the topology whose open sets are the intervals $[0, \alpha)$ ($\alpha \leq \mu_\beta$).*
- (6) *The topological space $\prod_{\beta \in \kappa} \mu_\beta$ does not satisfies CAP_λ^* , where each μ_β is endowed with the order topology.*

If $\lambda \geq \mu_\beta$, for every $\beta \in \kappa$, and $\kappa \geq \text{cf } S_\lambda(\lambda)$, then the preceding conditions are also equivalent to the following ones:

- (7) *The model $\langle \lambda, <, R_v, \gamma \rangle_{v \in V, \gamma < \lambda}$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{A}$ in a language with at most κ new symbols (and sorts) such that whenever $\mathfrak{B} \equiv \mathfrak{A}$ and $\mathfrak{B}$ (respectively, $\lambda^\mathfrak{B}$) has an element x such that $\mathfrak{B} \models \neg R_v(x)$ for every $v \in V$, then there exists $\beta \in \kappa$ such that $\mathfrak{B}$ has a μ_β -non standard element.*

- (8) *Every κ -(λ)-compact logic (equivalently, every κ -(λ)-compact logic generated by $\sup_{\beta \in \kappa} \mu_\beta$ cardinality quantifiers) is κ -(μ_β, μ_β)-compact for some $\beta \in \kappa$.*

If in addition λ is a regular cardinal, then the preceding conditions are also equivalent to the following one:

- (9) *The model $\langle \lambda, <, \gamma \rangle_{\gamma < \lambda}$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{A}$ in a language with at most κ new symbols (and sorts) such that whenever $\mathfrak{B} \equiv \mathfrak{A}$ and $\mathfrak{B}$ (respectively, $\lambda^{\mathfrak{B}}$) has a λ -non standard element, then, for some $\beta \in \kappa$, $\mathfrak{B}$ has a μ_β -non standard element.*

Proof. The equivalence of Conditions (1)-(3) is proved as in Part II, Theorem 1, equivalence of Conditions (b), (b'), (c).

(1) $\Rightarrow$ (4). Suppose by contradiction that (1) holds and (4) fails. Thus, there are topological spaces $(X_\beta)_{\beta \in \kappa}$ such that $X = \prod_{\beta \in \kappa} X_\beta$ satisfies CAP_λ^* , but, for every $\beta \in \kappa$, X_β fails to satisfy $\text{CAP}_{\mu_\beta}^*$.

This means that for every $\beta \in \kappa$ there exists a sequence $((y_\beta)_\alpha)_{\alpha < \mu_\beta}$ which has no μ_β -accumulation point in X_β .

Suppose that $(f_\beta)_{\beta \in \kappa}$ are functions as given by (1). Define a λ -indexed sequence $(x_\gamma)_{\gamma < \lambda}$ in $X = \prod_{\beta \in \kappa} X_\beta$, as follows. For $\gamma < \lambda$, $x_\gamma = \prod_{\beta \in \kappa} x_{\gamma, \beta}$ with $x_{\gamma, \beta} = (y_\beta)_{f_\beta(\gamma)}$.

Since X satisfies CAP_λ^* , the sequence $(x_\gamma)_{\gamma < \lambda}$ has a λ -accumulation point $x \in X$. By Proposition 2, there exists an ultrafilter D uniform over λ such that $(x_\gamma)_{\gamma < \lambda}$ D -converges to x .

By Lemma 5, for every $\beta \in \kappa$, the sequence $(x_{\gamma, \beta})_{\gamma < \lambda}$ D -converges in X_β . Since $x_{\gamma, \beta} = (y_\beta)_{f_\beta(\gamma)}$, we have that, for every $\beta \in \kappa$, the sequence $((y_\beta)_\alpha)_{\alpha < \mu_\beta}$ $f_\beta(D)$ -converges to some point of X_β .

By (1), there exists some $\bar{\beta} \in \kappa$ such that $f_{\bar{\beta}}(D)$ is uniform over $\mu_{\bar{\beta}}$, but then Proposition 2 implies that $((y_{\bar{\beta}})_\alpha)_{\alpha < \mu_{\bar{\beta}}}$ has a $\mu_{\bar{\beta}}$ -accumulation point, contradiction.

(4) $\Rightarrow$ (5) is trivial, since μ_β does not satisfies $\text{CAP}_{\mu_\beta}^*$, recalling the assumption that each μ_β is a regular cardinal.

(5) $\Rightarrow$ (6) is trivial, since the topology in (6) is finer than the topology in (5).

(6) $\Rightarrow$ (1). If (6) holds, then there exists a λ -indexed sequence $(x_\gamma)_{\gamma < \lambda}$ in X having no λ -accumulation point in X . Say, $x_\gamma = \prod_{\beta \in \kappa} (x_\gamma)_\beta$.

For $\beta \in \kappa$ and $\gamma < \lambda$, define $f_\beta(\gamma) = (x_\gamma)_\beta$. The contrapositive of Proposition 6 (a) $\Rightarrow$ (b) then implies (1).

The equivalence of (1), (7) and (9) is proved as in Part IV, Theorem 7 (Cf. also Part V, Theorem 1.2, Conditions (a), (d), (e)).

The equivalence of Condition (8) with the other conditions shall be presented elsewhere. $\square$

Proposition 9. *Suppose that Condition (4) in Theorem 8 holds. Then the following holds.*

If $(Y_j)_{j \in J}$ is a family of topological spaces, $\prod_{j \in J} Y_j$ satisfies CAP_λ^ and if, for every $\beta \in \kappa$, we put $J_\beta = \{j \in J \mid Y_j \text{ does not satisfy } \text{CAP}_{\mu_\beta}^*\}$, then there is $\beta \in \kappa$ such that $|J_\beta| \leq |\beta|$.*

Proof. Suppose that Condition (4) in Theorem 8 holds, but the conclusion of Proposition 9 fails. Thus, there exists a product $Y = \prod_{j \in J} Y_j$ satisfying CAP_λ^* , but such that $|J_\beta| > |\beta|$, for every $\beta \in \kappa$. Hence we can inductively construct a sequence $(j_\beta)_{\beta < \kappa}$ such that, for every $\beta < \kappa$, Y_{j_β} fails to satisfy $\text{CAP}_{\mu_\beta}^*$ and, moreover, for $\beta < \beta' < \kappa$, $j_\beta \neq j_{\beta'}$.

For $\beta \in \kappa$, put $X_\beta = Y_{j_\beta}$. Hence, $\prod_{\beta \in \kappa} X_\beta$ satisfies CAP_λ^* , since we are taking the product of a subset of the factors of Y . But then (4) implies that some $X_\beta = Y_{j_\beta}$ satisfies $\text{CAP}_{\mu_\beta}^*$, a contradiction. $\square$

We have a partial version of Theorem 8 for the case when the μ_β 's are not necessarily regular.

As far as Conditions (4)(5) below are concerned, for each $\beta \in \kappa$, choose some set $V_\beta \subseteq S_{\mu_\beta}(\mu_\beta)$ cofinal in $S_{\mu_\beta}(\mu_\beta)$ of cardinality $\leq \kappa$ and, for $v \in V_\beta$, let R_v be the unary predicate on μ_β defined by $R_v(\alpha)$ if and only if $\alpha \in v$.

Theorem 10. *Suppose that κ is a cardinal, λ is an infinite cardinal, and $(\mu_\beta)_{\beta \in \kappa}$ is a set of infinite cardinals. Then the following conditions are equivalent.*

- (1) *There are κ functions $(f_\beta)_{\beta \in \kappa}$ such that*
 - (a) *For each $\beta \in \kappa$, $f_\beta : \lambda \rightarrow \mu_\beta$; and*
 - (b) *for every ultrafilter D uniform over λ there is $\beta \in \kappa$ such that $f_\beta(D)$ is uniform over μ_β .*
- (2) *There are κ functions $(f_\beta)_{\beta \in \kappa}$ such that*
 - (a) *For each $\beta \in \kappa$, $f_\beta : \lambda \rightarrow \mu_\beta$; and*
 - (b) *for every function $g : \in \prod_{\beta \in \kappa} S_{\mu_\beta}(\mu_\beta)$ there exists some finite set $F \subseteq \kappa$ such that $\left| \bigcap_{\beta \in F} f_\beta^{-1}(g(\beta)) \right| < \lambda$.*
- (3) *There is a family $(C_{\alpha, \beta})_{\beta \in \kappa, \alpha \in \mu_\beta}$ of subsets of λ such that:*
 - (a) *For every $\beta \in \kappa$, $(C_{\alpha, \beta})_{\alpha \in \mu_\beta}$ is a partition of λ .*
 - (b) *For every function $g \in \prod_{\beta \in \kappa} S_{\mu_\beta}(\mu_\beta)$ there exists a finite subset $F \subseteq \kappa$ such that $\left| \bigcap_{\beta \in F} \bigcup_{\alpha \in g(\beta)} C_{\alpha, \beta} \right| < \lambda$.*

If $\kappa \geq \text{cf } S_\lambda(\lambda)$, and $\lambda \geq \mu_\beta$, $\kappa \geq \text{cf } S_{\mu_\beta}(\mu_\beta)$, for every $\beta \in \kappa$, then the preceding conditions are also equivalent to the following ones:

- (4) The model $\langle \lambda, <, R_v, \gamma \rangle_{v \in V \cup V_\beta, \gamma < \lambda}$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{A}$ in a language with at most κ new symbols (and sorts) such that whenever $\mathfrak{B} \equiv \mathfrak{A}$ and $\lambda^\mathfrak{B}$ has an element x such that $\mathfrak{B} \models \neg R_v(x)$ for every $v \in V$, then there exists $\beta \in \kappa$ such that $\mathfrak{B}$ has an element y in $\mu_\beta^\mathfrak{B}$ such that $\mathfrak{B} \models \neg R_v(y)$ for every $v \in V_\beta$.

If in addition λ is a regular cardinal, then the preceding conditions are also equivalent to the following one:

- (5) The model $\langle \lambda, <, R_v, \gamma \rangle_{v \in V_\beta, \gamma < \lambda}$ has an expansion (equivalently, a multi-sorted expansion) $\mathfrak{A}$ in a language with at most κ new symbols (and sorts) such that whenever $\mathfrak{B} \equiv \mathfrak{A}$ and $\mathfrak{B}$ has a λ -non standard element, then there exists $\beta \in \kappa$ such that $\mathfrak{B}$ has an element y in $\mu_\beta^\mathfrak{B}$ such that $\mathfrak{B} \models \neg R_v(y)$ for every $v \in V_\beta$.

Remark 11. Of course, there is the possibility of proving a mix between Theorems 8 and 10, in the case when certain μ_β 's are regular and other μ_β 's are singular.

We leave details to the reader.

Problems 12. (a) We do not know whether we can extend Theorem 10, for the case when the μ_β 's are singular, by adding further equivalent conditions analogue to Conditions (4)-(6) and (8) in Theorem 8.

(b) Does $\lambda \xrightarrow{\lambda^+} \mu$ implies $\lambda \xrightarrow{2^\lambda} \mu$?

(c) Find conditions equivalent to the conditions in Theorems 8 and 10 which are expressed in terms of Boolean Algebras.

Remark 13. In most cases, in our proofs, we are not using the full axiom of choice, but only the Prime Ideal Theorem.

REFERENCES

- [AU] P. Alexandroff, P. Urysohn, *Mémoire sur les espaces topologiques compacts*, Ver. Akad. Wetensch. Amsterdam **14** (1929), 1-96.
- [BF] J. Barwise, S. Feferman (editors), *Model-theoretic logics*. Perspectives in Mathematical Logic. Springer-Verlag, New York, 1985.
- [C1] X. Caicedo, *On productive $[\kappa, \lambda]$ -compactness, or the Abstract Compactness Theorem revisited*, manuscript (1995).
- [C2] X. Caicedo, *The Abstract Compactness Theorem Revisited*, in *Logic and Foundations of Mathematics (A. Cantini et al. editors)*, Kluwer Academic Publishers (1999), 131-141.
- [Ch] Chang, C. C. *Descendingly incomplete ultrafilters*. Trans. Amer. Math. Soc. **126** (1967) 108-118.
- [CK] C. C. Chang and J. Keisler, *Model Theory*, Amsterdam (1977).
- [CN] W. Comfort, S. Negrepointis, *The Theory of Ultrafilters*, Berlin (1974).

- [EU] P. Erdős, S. Ulam, *On equations with sets as unknowns*, Proc. Nat. Acad. Sci. U.S.A. **60** (1968), 1189–1195.
- [KM] A. Kanamori and M. Magidor, The evolution of large cardinal axioms in Set Theory, in: Higher Set Theory, edited by G. H. Müller and D. S. Scott, 99–275, Berlin (1978).
- [KV] K. Kunen and J. E. Vaughan (editors), *Handbook of Set Theoretical Topology*, Amsterdam (1984).
- [L1] P. Lipparini, *About some generalizations of (λ, μ) -compactness*, Proceedings of the 5th Easter conference on model theory (Wendisch Rietz, 1985), Seminarber., Humboldt-Univ. Berlin, Sekt. Math. **93**, 139–141 (1987). Available also at the author's web page.
- [L2] P. Lipparini, *Limit ultrapowers and abstract logics*, J. Symbolic Logic **52** (1987), no. 2, 437–454.
- [L3] P. Lipparini, *The compactness spectrum of abstract logics, large cardinals and combinatorial principles*, Boll. Unione Matematica Italiana ser. VII, **4-B** 875–903 (1990).
- [L4] P. Lipparini, *Ultrafilter translations, I: (λ, λ) -compactness of logics with a cardinality quantifier*, Arch. Math. Logic **35**, 63–87 (1996).
- [L5] P. Lipparini, *Productive $[\lambda, \mu]$ -compactness and regular ultrafilters*, Topology Proceedings **21** (1996), 161–171.
- [L6] P. Lipparini, *Compact factors in finally compact products of topological spaces*, Topology and its Applications, **153** (2006), 1365–1382.
- [L7] P. Lipparini, *Combinatorial and model-theoretical principles related to regularity of ultrafilters and compactness of topological spaces. I*, arXiv:0803.3498; *II*:0804.1445; *III*:0804.3737; *IV*:0805.1548 (2008); *V*:0903.4691 (2009).
- [S] Saks, Victor, Ultrafilter invariants in topological spaces, Trans. Amer. Math. Soc. **241** (1978), 79–97.
- [V1] J. E. Vaughan, *Some recent results in the theory of $[a, b]$ -compactness*, in *TOPO 72—General Topology and its Applications* (Proc. Second Pittsburg Internat. Conf., Carnegie-Mellon Univ. and Univ. Pittsburg, 1972), Lecture Notes in Mathematics **378** (1974), 534–550.

DIPARTITO DI MATEMATICA, VIALE DELLA RICERCA SCIENTIFICA, II UNIVERSITÀ ROMA (TOR VERGATA), I-00133 ROME ITALY
 URL: <http://www.mat.uniroma2.it/~lipparin>